# Selection of diploid and tetraploid banana hybrids resistant to *Pseudocercospora fijiensis*





Selection of Diploid and Tetraploid Banana Hybrids Resistant to *Pseudocercospora fijiensis* 

MDPI

Zalmar Santana Gonçalves<sup>1</sup>, Fernando Haddad<sup>2</sup>, Vanusia Batista de Oliveira Amorim<sup>2</sup>, Claudia Fortes Ferreira<sup>2</sup>, Saulo <sup>Zalmar Santana Gonçalves <sup>1,4</sup>, Anelita de Jesus Rocha<sup>1,4</sup>, Fernando Haddad<sup>2</sup>, Vanusia Batista de Oliveira Amorim<sup>2</sup>, Claudia Fortes Ferreira<sup>2</sup>, Saulo <sup>Almar Santana Gonçalves <sup>1,4</sup>, Anelita de Jesus Rocha<sup>1,4</sup>, Fernando Haddad<sup>2</sup>, Vanusia Batista de Oliveira Amorin<sup>2</sup>, Claudia Fortes Ferreira<sup>2</sup>, Saulo <sup>Almar Santana Gonçalves <sup>1,4</sup>, Anelita de Jesus Rocha<sup>1,4</sup>, Fernando Haddad<sup>2</sup>, Vanusia Batista de Oliveira Amorin<sup>2</sup>, Claudia Fortes Ferreira<sup>2</sup>, Saulo <sup>Almar Santana Gonçalves <sup>1,4</sup>, Anelita de Jesus Rocha<sup>1,4</sup>, Fernando Haddad<sup>2</sup>, Vanusia Batista de Oliveira Amorin<sup>2</sup>, Claudia Fortes Ferreira<sup>2</sup>, Saulo <sup>Almar Santana Gonçalves <sup>1,4</sup>, Anelita de Jesus Rocha<sup>1,4</sup>, Fernando Haddad<sup>2</sup>, Vanusia Batista de Oliveira Amorin<sup>2</sup>, Claudia Fortes Ferreira<sup>2</sup>, Saulo <sup>Almar Santana Gonçalves <sup>1,4</sup>, Anelita de Jesus Rocha<sup>1,4</sup>, Fernando Haddad<sup>2</sup>, Vanusia Batista de Oliveira Amorin<sup>2</sup>, Claudia Fortes Ferreira<sup>2</sup>, Saulo <sup>Almar Santana Gonçalves <sup>1,4</sup>, Anelita de Jesus Rocha<sup>1,4</sup>, Fernando Haddad<sup>2</sup>, Vanusia Batista de Oliveira Amorin<sup>2</sup>, Claudia Fortes Ferreira<sup>2</sup>, Saulo <sup>Almar Santana Gonçalves <sup>1,4</sup>, Anelita de Jesus Rocha<sup>1,4</sup>, Fernando Haddad<sup>2</sup>, Vanusia Batista de Oliveira Amorin<sup>2</sup>, Claudia Fortes Ferreira<sup>2</sup>, Almar Santa Gonçalves <sup>1,4</sup>, Anelita de Jesus Rocha<sup>1,4</sup>, Fernando Haddad<sup>2</sup>, Vanusia Batista de Oliveira Amorin<sup>2</sup>, Claudia Fortes Ferreira<sup>2</sup>, Almar Santa Gonçalves <sup>1,4</sup>, Anelita de Jesus Rocha<sup>1,4</sup>, Ferrando Haddad<sup>2</sup>, Vanusia Batista de Oliveira Amorin<sup>2</sup>, Claudia Fortes Ferreira<sup>2</sup>, Saulo <sup>1,4</sup>, Santa <sup>1,4</sup>, Santa <sup>1,4</sup>, Santa <sup>1,4</sup>, Ferrando Haddad<sup>2</sup>, <sup>1,4</sup>, <sup>1,4</sup>,</sup></sup></sup></sup></sup></sup></sup></sup>

<sup>1</sup>Universidade Estatual de Feira de Santana, Av. Transnordestina, s/n, Novo Horizonte, CEP 44036-900, Feira de Santana, Bahia, Brazil. E-mail: zalmarufrb@hotmail.com <sup>2</sup>Embrapa Mandioca e Fruticultura, Rua Embrapa, s/n CP 007, Chapadinha, Cruz das Almas, Bahia, Brazil. E-mails: fernando.haddad@embrapa.br; vanusiaamorim50@gmail.com; claudia.ferreira@embrapa.br, saulo.oliveira@embrapa.br.

\*Corresponding author: edson.amorim@embrapa.br. Telephone: +55 75 3312 8058.

#### Introduction, context and objectives

Bananas and plantains are cultivated in tropical and subtropical regions of more than 150 countries, on approximately 11 million hectares. Black Sigatoka can lead to 100% production loss in the absence of chemical control, which can increase the production cost by 30%, impact health and contaminate the environment. Developing cultivars resistant to black Sigatoka is the focus of many banana genetic breeding programs in the world and is an alternative to using fungicides. Thus, the objective of this work was to evaluate 23 diploid and eight tetraploid banana hybrids in the presence of *Pseudocercospora fijiensis*.

## Mat & Methods

The experiment was conducted in the experimental area of Embrapa in Brazil. The experimental design was completely randomized, with 10 replicates per genotype. Four selection strategies for genotypes resistant to black Sigatoka were evaluated: (1) evaluation of symptoms of the disease during the entire period of the experiment; (2) evaluation of symptoms during the summer months; (3) evaluation of symptoms during the winter months; and (4) classical evaluation of symptoms at the emission and harvest of bunches. The genotypes were evaluated for their behavior in the presence of black Sigatoka using a scale of scores of symptoms of the disease in the leaves, which was proposed by Stover (1972) and modified by Gauhl (1989). The evaluations started in the sixth month after planting when symptoms of black Sigatoka on leaf number 3 were observed until senescence. The scale of scores was applied at intervals of 15 days for the n<sup>o</sup> 3 leaves selected. The scores were recorded at intervals of 15 days and transformed into a disease severity index (DI, %) based on the formula described by Mckinney (1923). The area under the disease progress curve (AUDPC) was estimated with the formula proposed by Madden et al., (2007). Evaluations to select the hybrids in the summer (December to March) and winter (June to September) and estimates of the DI and AUDPC for these intervals (seasons of the year), were based on evaluations of the symptoms at 15-day intervals and the number of replicates per genotype. Based on the averages of the four DI and AUDPCs, a heat map analysis was conducted that produced a graphic interpretation where data referring to each genotypes. Using the same DIs and AUDPCs, a heat map analysis was conducted that produced a graphic interpretation where data referring to each genotypes are represented by colors; shades in the scale of green are associated with levels of genetic resistance and those in the scale of red indicate levels of susceptibility.

#### Results

Different groups formed depending on the evaluation strategy of the symptoms of black Sigatoka for both the number of genotypes and individuals in each cluster (Figure 1). From the dendrograms, it can be inferred that the **most efficient way** of **differentiating the genotypes** in the **resistant**, moderately resistant and **susceptible** classes is to **evaluate black Sigatoka symptoms** in the leaves during the **winter**. **Higher** mean **values** were observed in the **DI** and **AUDPC** for the evaluations made in the **winter**, **reinforcing** that this is the **best period** to **select genotypes resistant** to the disease, since there were better environmental conditions for the development of the epidemic in the field. With the goal of corroborating the cluster results described above, a **heat map** analysis was conducted based on the **four strategies** used to select the genotypes resistant to black Sigatoka (Figure 2). There was noticeable distinction among the selection periods based on the intensity of the colors in the figures where shades in the **green scale** are associated with levels of genetic **resistance** and those in the **red** scale indicate levels of **susceptibility**. For the set of improved diploids, the DI varied from 0.00 to 48.81 and the AUDPC varied from 0.00 to 2439.51. **Three** improved **diploids** were **completely resistant to black Sigatoka**: **CNPMF0519**, **CNPMF 0557 and CNPMF0612**. Another **nine** exhibited **quantitative resistance** to the pathogen, notably **CNPMF0811** and **CNPMF0993** that had the lowest DI and AUDPC among the genotypes. These five **genotypes** have potential for **use in crosses** with susceptible cultivars with the goal of **developing commercial cultivars resistant** to black Sigatoka. For the tetraploid hybrids evaluated for resistance to black Sigatoka, the DI varied from 15.06 and 63.52 and the AUDPC varied from 1000.21 to 3717.71. The genotype **CNPMF0906** (Prata type) and the cultivar BRS Princesa (Silk type) **exhibited moderate resistance** to the pathogen.

## **Conclusions and perspectives**

In the context of global **climate change**, according to probabilistic models associated with an increase in the average temperature, a 50% increase is estimated in areas on the planet cultivated with bananas, especially in subtropical regions, such as southern Brazil, and areas at higher elevations. In this scenario of **higher temperatures**, water will be supplied by irrigation, which will also **increase the severity and incidence of black Sigatoka**, since the fungal spores will be in more favorable conditions for germination (Calberto et al., 2015). Therefore, **cultivars adapted to this new condition** will be in **increasing demand** by farmers, especially those associated with genetic resistance to pests and diseases.





<u>References:</u> Stover, R. H. 1972. Banana Plantain and Abaca Diseases. Common Welth Mycological Institute, p.316.; Gauhl, F., Pasberg-Gauhl, C., Jones, D. R. 2000. Black leaf streak: disease cycle and epidemiology. In: Diseases of Banana, Abacá and Enset (Jones, D. R.), 2000, p.56-62.; Mckinney, H. H. 1923. Influence of soil, temperature and moisture on infection of wheat seedlings by Helminthosporium sativum. Journal of Agricultural Research, 26, 195-217.; Madden, L. V., Hugles, G., Bosch, F. V. D. 2007. The study of Plant Disease Epidemics. The American Phytopathologica Society, 32, 31-63. <u>Acknowledgements</u> : The National Council for Scientific and Technological Development (CNPq), the Fundação de Amparo à Pesquisa do Estado da Bahia (FAPESB), the Coordination for the Improvement of Higher Education Personnel (CAPES), the Bill & Melinda Gates Foundation and Embrapa.